

THE DUAL HILBERT-SAMUEL FUNCTION OF A MAXIMAL COHEN-MACAULAY MODULE

TONY J. PUTHENPURAKAL AND FAHED ZULFEQARR

ABSTRACT. Let R be a Cohen-Macaulay local ring with a canonical module ω_R . Let I be an $\mathfrak{m}$ -primary ideal of R and M , a maximal Cohen-Macaulay R -module. We call the function $n \mapsto \ell(\operatorname{Hom}_R(M, \omega_R/I^{n+1}\omega_R))$ the *dual Hilbert-Samuel function of M with respect to I* . By a result of Theodorescu this function is a polynomial function. We study its first two normalized coefficients.

INTRODUCTION

Let $(R, \mathfrak{m})$ be a Cohen-Macaulay local ring of dimension d with a canonical module ω_R . If the ring R is clear from context we write ω_R as ω . Let I be an $\mathfrak{m}$ -primary ideal and M , a finitely generated R -module.

The Hilbert-Samuel function of M with respect to I is the function

$$n \mapsto \ell\left(M \otimes \frac{R}{I^{n+1}}\right) \quad \text{for all } n \geq 0.$$

It is well-known (see [3, 4.6.2]) that for all large values of n , this function is given by a polynomial of degree equal to $\dim M$. This polynomial is called the *Hilbert-Samuel polynomial of M with respect to I* . Its normalized leading coefficient $e_0^I(M)$ is called the *multiplicity of M with respect to I* .

When M is Cohen-Macaulay R -module of dimension r , we define the *dual Hilbert-Samuel function with respect to I* by

$$n \mapsto \ell\left(\operatorname{Ext}_R^{d-r}(M, \frac{\omega}{I^{n+1}\omega})\right) \quad \text{for all } n \geq 0,$$

We call the power-series

$$D(M, t) = \sum_{n \geq 0} \ell\left(\operatorname{Ext}_R^{d-r}(M, \frac{\omega}{I^{n+1}\omega})\right) z^n$$

as the *dual Hilbert-Samuel series of M with respect to I* . We concentrate on the case when $\dim M = \dim R$, i.e., when M is a maximal Cohen-Macaulay (MCM) R -module.

For $i \geq 0$ we call the function

$$n \mapsto \ell\left(\operatorname{Ext}_R^i(M, \frac{\omega}{I^{n+1}\omega})\right) \quad \text{for all } n \geq 0$$

as the *i -th dual Hilbert-Samuel function of M with respect to I* . We study these functions for $i = 0, 1$ and when M is MCM. Using Theodorescu's result [9, 4], it follows that this function is given by a polynomial in n for all large values of

Date: September 19, 2008.

1991 Mathematics Subject Classification. Primary 13D45 ; Secondary 13A30.

Key words and phrases. multiplicity, reduction, Hilbert-Samuel polynomial .

n . We denote this polynomial by $\varepsilon_M^i(I, t)$ and call it the i -th dual Hilbert-Samuel polynomial of M with respect to I . Furthermore we have

- (1) for $i = 0$, the dual Hilbert-Samuel polynomial $\varepsilon_M^0(I, t)$ is of degree d .
- (2) for each $i > 0$, the i -th dual Hilbert-Samuel polynomial $\varepsilon_M^i(I, t)$ is of degree at most $d - 1$.

Therefore the polynomial $\varepsilon_M^0(I, t)$ can be written in the following form

$$\varepsilon_M^0(I, t) = \sum_{i=0}^d (-1)^i c_i^I(M) \binom{t+d-i}{d-i}.$$

We call the integers $c_0^I(M), c_1^I(M), \dots, c_d^I(M)$ the dual Hilbert coefficients of M with respect to I . In Proposition 2.5 we show that

$$c_0^I(M) = e_0^I(M).$$

This also shows that if J is a reduction of I with respect to M then $c_0^J(M) = c_0^I(M)$.

One might also try to plausibly study the "Dual Hilbert function" of a MCM module M , i.e., the function

$$\delta_I(M, n) = \ell(\operatorname{Hom}_R(M, I^n \omega / I^{n+1} \omega)) \quad \text{for } n \geq 0.$$

However we feel that it is not as interesting as the Dual Hilbert-Samuel function for the following reasons: Assume R is Gorenstein and $I = \mathfrak{m}$.

- The dual Hilbert function is

$$\delta_{\mathfrak{m}}(M, n) = \mu(M) \ell(\mathfrak{m}^n / \mathfrak{m}^{n+1}).$$

Here $\mu(M)$ denotes the number of minimal generators of M . Thus this function tells very little information on M .

- If J is a minimal reduction of $\mathfrak{m}$ with respect to M then it can be easily shown that the normalized leading coefficient of the polynomial function $\delta_J(M, n)$ is $e_0^J(M) = e_0^{\mathfrak{m}}(M)$. Using the above calculation we get that the normalized leading coefficient of $\delta_{\mathfrak{m}}(M, n)$ is $\mu(M)e_0^{\mathfrak{m}}(A)$. Since M is MCM we get that $e_0^{\mathfrak{m}}(M) = \mu(M)e_0^{\mathfrak{m}}(A)$ if and only if M is free. Thus for interesting cases we do have that leading coefficients of $\delta_J(M, n)$ and $\delta_{\mathfrak{m}}(M, n)$ are different.

A crucial property of the dual Hilbert-Samuel function is that it behaves well modulo a sufficiently general element in $I \setminus \mathfrak{m}I$. More specifically in Proposition 3.3 we show the following: Let $x \in I \setminus \mathfrak{m}I$ be sufficiently general. Set $S = R/xR$, $J = I/(x)$ and $N = M/xM$. We prove

$$c_i^J(N) = c_i^I(N) \quad \text{for } i = 0, \dots, d-1.$$

In Theorem 4.1 we show that if M is not a free MCM R -module then the function

$$(*) \quad n \longmapsto \ell\left(\operatorname{Ext}_R^1(M, \frac{\omega}{\mathfrak{m}^{n+1}\omega})\right) \quad \text{for } n \geq 0$$

is a polynomial type of degree $d-1$. Example 2.6 shows that $\deg \varepsilon_M^1(I, t)$ can be $-\infty$ in general. A consequence of $(*)$ is that if M is a non-free MCM R -module then

$$\mu(M)e_1(\omega_A) > c_1(M) + c_1(\operatorname{Syz}_1^R(M)).$$

In case the ring R is Gorenstein and its associated ring $G_I(R)$ is Cohen-Macaulay, we prove (see Theorem 5.7) that if r is a reduction number of I then we have

$$c_1^I(M) \geq r \cdot e_0^I(M) - \sum_{j=0}^d \sum_{n=0}^{r-1} \binom{d}{j} \ell \left(\text{Ext}_R^j(M, \frac{R}{I^{n-j+1}}) \right).$$

In view of the above result we study the function

$$\Phi^I(M) = \sum_{j=0}^d \sum_{n=0}^{r-1} \binom{d}{j} \ell \left(\text{Ext}_R^j(M, \frac{R}{I^{n+1-j}}) \right).$$

If $x \in I \setminus \mathfrak{m}I$ is an R -superficial element then we prove

$$\Phi^I(M) \geq \Phi^{I/(x)}(M/xM).$$

The dual Hilbert-Samuel function behaves well with respect to the basic properties just as the usual Hilbert-Samuel function. However for computational reasons it compares poorly with respect to the usual Hilbert function. In section 6 we compute the dual Hilbert-Samuel series of an Ulrich module M over a Gorenstein local ring R such that $G_{\mathfrak{m}}(R)$ is also Gorenstein. This computation is modulo a result from [5].

We now describe in brief the contents of this paper. In section 1 we introduce notation and discuss a few preliminary facts that we need. In section 2 we prove that $c_0^I(M) = e_0^I(M)$. In section 3 we discuss the behavior of the functions $\varepsilon^0, \varepsilon^1$ modulo a general element in $I \setminus \mathfrak{m}I$. In section 4 we prove that the polynomial $\varepsilon_M^1(\mathfrak{m}, t)$ is either zero or a polynomial of degree $d-1$. In section 5 we discuss a lower bound on $c_1^I(M)$. Finally in the last section we compute the dual Hilbert-Samuel series of an Ulrich module M over a Gorenstein local ring R such that $G_{\mathfrak{m}}(R)$ is also Gorenstein.

1. NOTATION AND PRELIMINARIES

Throughout this paper unless otherwise stated we assume that R is a Cohen-Macaulay (CM) local ring of dimension d with maximal ideal $\mathfrak{m}$ and that residue field $R/\mathfrak{m}$ is infinite. We also assume that R has a canonical module ω . Let I be an $\mathfrak{m}$ -primary ideal and M , a finitely generated R -module.

1.1. Notation : We denote the Rees Algebra of ideal I by $\mathcal{R}(I)$, the associated graded ring with respect to I by $G_I(R)$ and their associated graded modules by $\mathcal{R}(I, M)$ and $G_I(M)$ respectively, i.e.,

$$\begin{aligned} \mathcal{R}(I) &= \bigoplus_{n \geq 0} I^n, & G_I(R) &= \bigoplus_{n \geq 0} \frac{I^n}{I^{n+1}} \\ \text{and } \mathcal{R}(I, M) &= \bigoplus_{n \geq 0} I^n M, & G_I(M) &= \bigoplus_{n \geq 0} \frac{I^n M}{I^{n+1} M} \end{aligned}$$

Clearly $\mathcal{R}(I, M)$ is a finitely generated $\mathcal{R}(I)$ -module and $G_I(M)$ is a finitely generated $G_I(R)$ -module.

Since I is $\mathfrak{m}$ -primary ideal in R then by [3, 4.5.13], $l(I) = \dim R = d$. Using [9, 3(c)], since ω is faithful R -module, we get $l_{\omega}(I) = l(I)$. Thus $l_{\omega}(I) = d$.

Remarks 1.2. From [9, 4], it follows that the function

$$n \mapsto \ell \left(\text{Ext}_R^i(M, \frac{\omega}{I^{n+1}\omega}) \right) \quad \text{for each } i \geq 0$$

is given by a polynomial, say $\varepsilon_M^i(I, n)$, in n for all $n \gg 0$ and

$$(*) \quad \deg \varepsilon_M^i(I, t) \leq \max \{ \dim \text{Ext}_R^i(M, \omega), l_\omega(I) - 1 \}.$$

Moreover equality holds if $\dim \text{Ext}_R^i(M, \omega) \geq l(I) = d$. If M is an *MCM* R -module then using $(*)$ we get that the function

- (a) $n \mapsto \ell (\text{Hom}_R(M, \omega/I^{n+1}\omega))$ is a polynomial type of degree d .
- (b) $n \mapsto \ell (\text{Ext}_R^1(M, \omega/I^{n+1}\omega))$ is a polynomial type of degree at most $d - 1$.

2. RELATION WITH HILBERT COEFFICIENTS

In this section we establish an equality between $e_0^I(M)$ and the dual Hilbert coefficient $c_0^I(M)$ of M .

2.1. Notation: We set

$$\mathcal{F} \equiv \{M_n^\dagger\}_{n \geq 0} \quad \text{where} \quad M_n^\dagger = \text{Hom}_R(M, I^n \omega) \quad \text{for all } n \geq 1$$

$$E_I^i(M) = \bigoplus_{n \geq 0} \text{Ext}_R^i(M, I^n \omega) \quad \text{and} \quad D_I^i(M) = \bigoplus_{n \geq 0} \text{Ext}_R^i(M, \omega/I^n \omega).$$

Remarks 2.2.

- (a) It can be easily seen that $\mathcal{F}$ is an stable I -filtration. Therefore by [1, 11.4], the function

$$n \mapsto \ell (M^\dagger / M_n^\dagger)$$

is given by a polynomial of degree $\dim M^\dagger = d$, for all $n \gg 0$. Further the normalized leading coefficient of this function is $e_0^I(M^\dagger)$.

- (b) Notice that M and $\text{Hom}_R(M, \omega)$ are both *MCM* R -modules, It is well known and can be easily verified that

$$e_0^I(M^\dagger) = e_0^I(M).$$

- (c) Let $E = \bigoplus_{n \geq 0} E_n$ be a finitely generated $\mathcal{R}(I)$ -module such that $\ell_R(E_n) < \infty$ for all $n \geq 0$. It is well known that the function $n \mapsto \ell(E_n)$ is polynomial of degree $\leq d - 1$.

The following lemma is crucial.

Lemma 2.3. *Let M be a finitely generated R -module. Then for each $i \geq 0$, $E_I^i(M)$ is finitely generated graded $\mathcal{R}(I)$ -module.*

Proof. Consider a minimal free resolution of M

$$\mathbb{F}_\bullet : \dots \rightarrow F_2 \rightarrow F_1 \rightarrow F_0 \rightarrow 0.$$

Applying the functor $\text{Hom}_R(-, \bigoplus_{n \geq 0} I^n \omega)$ to this sequence, we get the complex $\text{Hom}_R(\mathbb{F}_\bullet, \bigoplus_{n \geq 0} I^n \omega)$ of finitely generated graded $\mathcal{R}(I)$ -modules. We now have

$$\begin{aligned} \bigoplus_{n \geq 0} \text{Ext}_R^i(M, I^n \omega) &\simeq \text{Ext}_R^i(M, \bigoplus_{n \geq 0} I^n \omega) \\ &= \mathcal{H}^i \left(\text{Hom}_R(\mathbb{F}_\bullet, \bigoplus_{n \geq 0} I^n \omega) \right). \end{aligned}$$

Hence for each $i \geq 0$, $\bigoplus_{n \geq 0} \text{Ext}_R^i(M, I^n \omega)$ is finitely generated graded $\mathcal{R}(I)$ -module. $\square$

Lemma 2.4. *Let M be an MCM R -module. Then the following hold :*

- (a) *For $i \geq 1$, $D_I^i(M) \cong E_I^{i+1}(M)$. So $D_I^i(M)$ is a finitely generated graded $\mathcal{R}(I)$ -module.*
- (b) *For all $i \geq 1$, the function $n \mapsto \ell(\text{Ext}_R^i(M, I^n \omega))$ is a polynomial function of degree $\leq d - 1$.*

Proof. (a): Consider the exact sequence

$$0 \rightarrow I^n \omega \rightarrow \omega \rightarrow \frac{\omega}{I^n \omega} \rightarrow 0.$$

Since M is MCM, $\text{Ext}_R^i(M, \omega) = 0$ for all $i \geq 1$. So applying the functor $\text{Hom}_R(M, -)$ to the above exact sequence we get

$$\text{Ext}_R^i(M, \omega/I^n \omega) \simeq \text{Ext}_R^{i+1}(M, I^n \omega) \quad \text{for all } i \geq 1.$$

From Lemma 2.3 it follows that $D_I^i(M)$ is finitely generated graded $\mathcal{R}(I)$ -module and $D_I^i(M) \cong E_I^{i+1}(M)$ for all $i \geq 1$.

(b): Since I is $\mathfrak{m}$ -primary and by using part (a), we get that for all $i \geq 2$, $\ell(\text{Ext}_R^i(M, I^n \omega)) < \infty$ for all n . Therefore by 2.2(c) the function $n \mapsto \ell(\text{Ext}_R^i(M, I^n \omega))$ is a polynomial function of degree at most $l(I) - 1 = d - 1$. For $i = 1$, consider the exact sequence

$$0 \rightarrow I^n \omega \rightarrow \omega \rightarrow \frac{\omega}{I^n \omega} \rightarrow 0.$$

Applying the functor $\text{Hom}_R(M, -)$ we get

$$0 \rightarrow \frac{M^\dagger}{M_n^\dagger} \rightarrow \text{Hom}_R(M, \frac{\omega}{I^n \omega}) \rightarrow \text{Ext}_R^1(M, I^n \omega) \rightarrow 0.$$

It follows that $\ell(\text{Ext}_R^1(M, I^n \omega)) < \infty$ for all $n \geq 1$. Using 2.2(c) we get that for all $n \gg 0$, the function $n \mapsto \ell(\text{Ext}_R^1(M, I^n \omega))$ is given a polynomial of degree at most $d - 1$. $\square$

Another useful observation is recorded in the following proposition.

Proposition 2.5. *The function $n \mapsto \ell(\text{Hom}_R(M, \omega_R/I^{n+1}\omega_R))$ is of polynomial type and the dual Hilbert coefficient $c_0^I(M)$ is same as the multiplicity of M with respect to I .*

Proof. Consider the exact sequence

$$0 \rightarrow I^n \omega \rightarrow \omega \rightarrow \frac{\omega}{I^n \omega} \rightarrow 0.$$

Applying the functor $\text{Hom}_R(M, -)$ we get

$$0 \rightarrow \frac{M^\dagger}{M_n^\dagger} \rightarrow \text{Hom}_R(M, \frac{\omega}{I^n \omega}) \rightarrow \text{Ext}_R^1(M, I^n \omega) \rightarrow 0.$$

Therefore it follows that

$$\ell\left(\text{Hom}_R(M, \frac{\omega}{I^n \omega})\right) = \ell\left(\frac{M^\dagger}{M_n^\dagger}\right) + \ell(\text{Ext}_R^1(M, I^n \omega)).$$

By Lemma 2.4(b), we have that the function $n \mapsto \ell(\text{Ext}_R^1(M, I^n \omega))$ is given a polynomial function of degree at most $d - 1$. Also by Remark 2.2(a), the

function $n \mapsto \ell(M^\dagger/M_n^\dagger)$ is a polynomial type of degree d . So the function $n \mapsto \ell(\text{Hom}_R(M, \omega_R/I^{n+1}\omega_R))$ is given a polynomial $\varepsilon_M^0(I, t)$ and the leading coefficient of this polynomial is same as that of $n \mapsto \ell(M^\dagger/M_n^\dagger)$. Therefore $c_I^0(M) = e_0^I(M^\dagger)$. Remark 2.2(b) gives that $c_0^I(M) = e_0^I(M)$. $\square$

We end this section by computing the dual Hilbert-Samuel function with respect to a parameter ideal J . The reasons for this computations are the following

- (1) If J is a minimal reduction of I with respect to $\mathfrak{m}$ then J is a parameter ideal. We also have

$$c_0^I(M) = e_0^I(M) = e_0^J(M) = c_0^J(M).$$

Here the first and the third equality is due to Proposition 2.5 while the second equality is a basic property of reductions.

- (2) The computation will also prove that

$$\text{Ext}_R^1\left(M, \frac{\omega}{J^{n+1}\omega}\right) = 0 \quad \text{for all } n \geq 0.$$

This assertion proves that we cannot hope that Theorem 4.2 to work for all $\mathfrak{m}$ -primary ideals.

- (3) We need to compute the Dual Hilbert-Samuel function in this essentially simplest case of an $\mathfrak{m}$ -primary ideal.

Example 2.6. Let M be a MCM module. Let $J = (x_1, \dots, x_d)$ be a parameter ideal. Then

- (a) $\varepsilon_M^0(J, n) = \ell\left(\text{Hom}_R(M, \frac{\omega}{J^{n+1}\omega})\right) = \ell(M/JM) \binom{n+d}{d}$
- (b) $\varepsilon_M^1(J, n) = \ell\left(\text{Ext}_R^1(M, \frac{\omega}{J^{n+1}\omega})\right) = 0 \quad \text{for all } n \geq 0.$

Proof. Let $\delta(n) = \mu(J^n) = \binom{n+d-1}{d-1}$. Since $x_1, \dots, x_d$ is a R -regular sequence, it is also ω -regular. So we get

$$\frac{J^n \omega}{J^{n+1} \omega} = \left(\frac{\omega}{J\omega}\right)^{\delta(n)}.$$

Set $S = R/J$. Notice $\omega_S = \omega/J\omega$. We will also use the fact that for $i \geq 0$,

$$\begin{aligned} \ell(\text{Ext}_R^i(M, \omega/J\omega)) &= \ell(\text{Ext}_S^i(M/JM, \omega_S)), \quad \text{cf. [4, §18, Lemma 2]} \\ (*) \quad &= \begin{cases} \ell(M/JM), & \text{if } i = 0; \\ 0, & \text{if } i \geq 1. \end{cases} \end{aligned}$$

We prove assertions (a), (b) by induction on n . For $n = 0$ the result follows from (*). For $n \geq 0$ we use the exact sequence

$$0 \longrightarrow \frac{J^n \omega}{J^{n+1} \omega} \longrightarrow \frac{\omega}{J^{n+1} \omega} \longrightarrow \frac{\omega}{J^n \omega} \longrightarrow 0.$$

We now apply the functor $\text{Hom}_R(M, -)$. The corresponding long exact sequence and (*) gives the result. $\square$

3. BEHAVIOR WITH RESPECT TO A SUPERFICIAL ELEMENTS

In this section we discuss the behavior of $\varepsilon_M^0(\omega, t)$ with respect to a superficial element. To ensure good behavior, see 3.3, it is important to choose superficial elements with some care, see 3.2.

Let us first recall some basic definitions.

3.1. An element $x \in I$ is called *M-superficial* with respect to I if there exists an integer $c \geq 0$ such that

$$(I^{n+1}M :_M x) \cap I^c M = I^n M \quad \text{for all } n \geq c.$$

Also recall that if $E = \bigoplus_{n \geq 0} E_n$ is finitely generated graded $\mathcal{R}(I)$ -module then an element $xt \in \mathcal{R}(I)_1$ is called *E-filter regular* if

$$(0 :_E x)_n = 0 \quad \text{for all } n \gg 0.$$

Equivalently $\alpha_x : E_n \rightarrow E_{n+1}$ are injective for $n \gg 0$,

where α_x is the map induced by multiplication by x (i.e., $\alpha_x(u) = xu$). Notice that if the residue field $R/\mathfrak{m}$ is infinite then superficial and filter regular element exists (see [2, 18.3.10]).

3.2. Choosing a superficial element: Since by Lemma 2.4, $D_I^i(M)$ are finitely generated $\mathcal{R}(I)$ -modules for $i \geq 1$, we can choose an element $x \in I \setminus I^2$ such that

- (a) x is I -superficial element with respect to $R \oplus \omega$.
- (b) $xt \in \mathcal{R}(I)_1$ is $D_I^1(M) \oplus D_I^2(M)$ -filter regular.

Clearly we have x is $R \oplus M$ -regular. Set $S = R/xR$, $J = I/(x)$ and $N = M/xM$. Notice that $\omega_S = \omega/x\omega$, the canonical module of S .

Proposition 3.3. *Let $x \in I \setminus I^2$ be an element satisfying 3.2. Then*

- (a) $\varepsilon_M^0(I, n) - \varepsilon_M^0(I, n-1) = \varepsilon_N^0(J, n)$ for $n \gg 0$.
- (b) $\varepsilon_M^1(I, n) - \varepsilon_M^1(I, n-1) = \varepsilon_N^1(J, n)$ for $n \gg 0$.

Proof. Consider the exact sequence

$$0 \rightarrow \frac{I^{n+1}\omega : x}{I^n\omega} \rightarrow \frac{\omega}{I^n\omega} \xrightarrow{\alpha_x^n} \frac{\omega}{I^{n+1}\omega} \rightarrow \frac{\omega}{(x, I^{n+1})\omega} \rightarrow 0,$$

such that $\alpha_x^n(u + I^n\omega) = xu + I^{n+1}\omega, \quad \forall u \in \omega.$

As x is ω -superficial and ω -regular, we have $I^{n+1}\omega : x = I^n\omega$ for $n \gg 0$. Set $J = I/(x)$. For $n \gg 0$, one can thus write above exact sequence as follows

$$0 \rightarrow \frac{\omega}{I^n\omega} \xrightarrow{\alpha_x^n} \frac{\omega}{I^{n+1}\omega} \rightarrow \frac{\omega_S}{J^{n+1}\omega_S} \rightarrow 0,$$

where $\omega_S = \omega_R/x\omega_R$, the canonical module of S . Now applying the functor $\text{Hom}_R(M, -)$ we get a long exact sequence

$$\begin{aligned} 0 \rightarrow \text{Hom}_R(M, \frac{\omega}{I^n\omega}) &\rightarrow \text{Hom}_R(M, \frac{\omega}{I^{n+1}\omega}) \rightarrow \text{Hom}_R(M, \frac{\omega_S}{J^{n+1}\omega_S}) \\ &\rightarrow \text{Ext}_R^1(M, \frac{\omega}{I^n\omega}) \rightarrow \text{Ext}_R^1(M, \frac{\omega}{I^{n+1}\omega}) \rightarrow \dots \end{aligned}$$

By our hypotheses on x , the maps

$$\begin{aligned} \text{Ext}_R^1(M, \alpha_x^n) : \text{Ext}_R^1(M, \frac{\omega}{I^n\omega}) &\rightarrow \text{Ext}_R^1(M, \frac{\omega}{I^{n+1}\omega}) \\ \text{Ext}_R^2(M, \alpha_x^n) : \text{Ext}_R^2(M, \frac{\omega}{I^n\omega}) &\rightarrow \text{Ext}_R^2(M, \frac{\omega}{I^{n+1}\omega}) \end{aligned}$$

are injective for all $n \gg 0$. Therefore for all $n \gg 0$, we get the exact sequences

$$\begin{aligned} 0 \rightarrow \operatorname{Hom}_R(M, \frac{\omega}{I^n \omega}) &\rightarrow \operatorname{Hom}_R(M, \frac{\omega}{I^{n+1} \omega}) \rightarrow \operatorname{Hom}_R(M, \frac{\omega_S}{J^{n+1} \omega_S}) \rightarrow 0 \\ 0 \rightarrow \operatorname{Ext}_R^1(M, \frac{\omega}{I^n \omega}) &\rightarrow \operatorname{Ext}_R^1(M, \frac{\omega}{I^{n+1} \omega}) \rightarrow \operatorname{Ext}_R^1(M, \frac{\omega_S}{J^{n+1} \omega_S}) \rightarrow 0. \end{aligned}$$

Using [4, Lemma 18.2], we get

$$\operatorname{Ext}_R^i(M, \frac{\omega_S}{J^{n+1} \omega_S}) \cong \operatorname{Ext}_S^i(N, \frac{\omega_S}{J^{n+1} \omega_S}) \quad \text{for all } n \geq 0.$$

The result follows. $\square$

The following is a very useful consequence to above proposition.

Corollary 3.4. *Let $x \in I \setminus I^2$ be an element satisfying 3.2. Then*

$$c_i^J(N) = c_i^I(M) \quad \text{for } i = 0, 1, 2, \dots, d-1.$$

Proof. It follows easily from Proposition 3.3(a) and the binomial identity

$$\binom{n+d-i-1}{d-i-1} = \binom{n+d-i}{d-i} - \binom{n+d-i-1}{d-i}.$$

$\square$

4. THE FUNCTION $n \mapsto \ell(\operatorname{Ext}_R^1(M, \omega/\mathfrak{m}^{n+1}\omega))$

In this section our goal is to show that for a non-free MCM R -module M of dimension $d \geq 1$, the degree of polynomial $\varepsilon_M^1(\mathfrak{m}, t)$ with respect to maximal ideal $\mathfrak{m}$ is exactly $d-1$. This need not be true for all $\mathfrak{m}$ -primary ideals in general (see Example 2.6).

Recall the i -th betti number β_i of M (see [3, 1.3.1]) is given by

$$\beta_i = \dim_k \operatorname{Ext}_R^i(M, k), \quad \text{where } k = R/\mathfrak{m}.$$

Note that $\beta_0 = \mu(M)$, the minimal number of generators of M .

The following is the dual version of Proposition 17 in [7].

Proposition 4.1. *Let $\dim M = d \geq 1$. Set $L = \operatorname{Syz}_R^1(M)$, the first syzygy of M . Then*

(a) *We have*

$$\sum_{n \geq 0} \ell(\operatorname{Ext}_R^1(M, \omega/I^{n+1}\omega)) t^n = \frac{f_M^I(t) - \mu(M)h_\omega^I(t) + f_L^I(t)}{(1-t)^{d+1}}.$$

(b) *For $n \gg 0$ the function $n \mapsto \ell(\operatorname{Ext}_R^1(M, \omega/I^{n+1}\omega))$ is given by a polynomial $\varepsilon_M^1(I, t)$ of the form*

$$\varepsilon_M^1(I, t) = (\mu(M) \cdot e_1^I(\omega) - c_1^I(M) - c_1^I(L)) \frac{t^{d-1}}{(d-1)!} + \text{lower degree terms}.$$

(c) $c_1^I(M) + c_1^I(L) \leq \mu(M) \cdot e_1^I(\omega)$.

Proof. (a): Consider the exact sequence

$$(i) \quad 0 \longrightarrow L \longrightarrow R^{\mu(M)} \longrightarrow M \longrightarrow 0.$$

Applying $\text{Hom}_R(-, \omega/I^{n+1}\omega)$ to this, we get

$$\begin{aligned} 0 \rightarrow \text{Hom}_R(M, \frac{\omega}{I^{n+1}\omega}) &\rightarrow \text{Hom}_R(R^{\mu(M)}, \frac{\omega}{I^{n+1}\omega}) \rightarrow \text{Hom}_R(L, \frac{\omega}{I^{n+1}\omega}) \\ &\rightarrow \text{Ext}_R^1(M, \frac{\omega}{I^{n+1}\omega}) \rightarrow 0. \end{aligned}$$

This induces that

$$\begin{aligned} \sum_{n \geq 0} \ell(\text{Ext}_R^1(M, \omega/I^{n+1}\omega)) t^n &= \frac{f_M^I(t)}{(1-t)^{d+1}} - \frac{\mu(M) \cdot h_\omega^I(t)}{(1-t)^{d+1}} + \frac{f_L^I(t)}{(1-t)^{d+1}} \\ &= \frac{f_M^I(t) - \mu(M) \cdot h_\omega^I(t) + f_L^I(t)}{(1-t)^{d+1}} \end{aligned}$$

(b): Now we set $p(t) = f_M^I(t) - \mu(M) \cdot h_\omega^I(t) + f_L^I(t)$. From Proposition 2.5, it follows that

$$p(1) = e_0^I(M) - \mu(M) \cdot e_0^I(\omega) + e_0^I(L) = 0, \quad \text{by (i).}$$

We thus write $p(t) = (1-t)s(t)$. This gives that

$$\sum_{n \geq 0} \ell(\text{Ext}_R^1(M, \omega/I^{n+1}\omega)) t^n = \frac{s(t)}{(1-t)^d}$$

This shows that $\deg \varepsilon_M^1(I, t) \leq d-1$. We also have

$$s(1) = -p'(1) = \mu(M) \cdot e_1^I(\omega) - c_1^I(M) - c_1^I(L).$$

Hence

$$\varepsilon_M^1(I, t) = (\mu(M) \cdot e_1^I(\omega) - c_1^I(M) - c_1^I(L)) \frac{t^{d-1}}{(d-1)!} + \text{lower terms in } t.$$

(c): This follows easily from (b). $\square$

In the following theorem we take $I = \mathfrak{m}$. We prove that if M is non-free MCM R -module then $\deg \varepsilon_M^1(\mathfrak{m}, t) = d-1$.

Theorem 4.2. *Let R be Cohen-Macaulay local of dimension $d \geq 1$. The following conditions are equivalent*

- (a) M is free.
- (b) $\varepsilon_M^1(\mathfrak{m}, t) = 0$.
- (c) $\deg \varepsilon_M^1(\mathfrak{m}, t) < d-1$.
- (d) $c_1(M) + c_1(\text{Syz}_R^1(M)) = \mu(M) \cdot e_1(\omega)$.

Example 2.6 shows that this result need not true for all m -primary ideals.

Proof of the theorem 4.2. The implications (a) $\Rightarrow$ (b) $\Rightarrow$ (c) are clear. The assertion (c) $\Rightarrow$ (d) follows from Proposition 4.1(b).

(d) $\Rightarrow$ (a): We apply induction on $d = \dim M$. If $d = 1$ then $\varepsilon_M^1(\mathfrak{m}, t) = 0$. So we get

$$\text{Ext}_R^1(M, \frac{\omega}{\mathfrak{m}^{n+1}\omega}) = 0 \quad \text{for all } n \gg 0.$$

The exact sequence

$$0 \longrightarrow \frac{\mathfrak{m}^n \omega}{\mathfrak{m}^{n+1} \omega} \longrightarrow \frac{\omega}{\mathfrak{m}^{n+1} \omega} \longrightarrow \frac{\omega}{\mathfrak{m}^n \omega} \longrightarrow 0$$

gives rise to a long exact sequence, for $n \gg 0$,

$$\begin{aligned} 0 \rightarrow \operatorname{Hom}_R(M, \frac{\mathfrak{m}^n \omega}{\mathfrak{m}^{n+1} \omega}) &\rightarrow \operatorname{Hom}_R(M, \frac{\omega}{\mathfrak{m}^{n+1} \omega}) \rightarrow \operatorname{Hom}_R(M, \frac{\omega}{\mathfrak{m}^n \omega}) \\ &\rightarrow \operatorname{Ext}_R^1(M, \frac{\mathfrak{m}^n \omega}{\mathfrak{m}^{n+1} \omega}) \rightarrow 0. \end{aligned}$$

Therefore we get a relation

$$\mu(M) \cdot e_0(\omega) - [e_0(M) \cdot n - c_1(M)] + [e_0(M) \cdot (n-1) - c_1(M)] - \beta_1 \cdot e_0(\omega) = 0.$$

(i) So, $e_0(M) = \mu(M) \cdot e_0(\omega) - \beta_1 \cdot e_0(\omega) = \mu(M) \cdot e_0(R) - \beta_1 \cdot e_0(R)$.

Set $L_1 = \operatorname{Syz}_R^1(M)$ and $L_2 = \operatorname{Syz}_R^1(L_1)$. As M is MCM so are L_1 and L_2 . The exact sequence

$$0 \rightarrow L_1 \rightarrow R^{\mu(M)} \rightarrow M \rightarrow 0$$

yields

$$(ii) \quad e_0(L_1) = \mu(M) \cdot e_0(R) - e_0(M).$$

Similarly we get

$$(iii) \quad e_0(L_2) = \beta_1 \cdot e_0(R) - e_0(L_1).$$

Therefore we have

$$\begin{aligned} e_0(L_2) &= \beta_1 \cdot e_0(R) - [\mu(M) \cdot e_0(R) - e_0(M)] \quad (\text{by (ii)}) \\ &= [\beta_1 \cdot e_0(R) - \mu(M) \cdot e_0(R)] + e_0(M) \\ &= -e_0(M) + e_0(M) = 0 \quad (\text{by (i)}). \end{aligned}$$

So $L_2 = 0$ and hence $\operatorname{projdim}_R(M) < \infty$. Using Auslander-Buchsbaum's formula we get M is free.

When $d > 1$. We choose $x_1, x_2, \dots, x_d \in \mathfrak{m} \setminus \mathfrak{m}^2$ such that

- (1) x_i is $R_{i-1} \oplus \omega_{i-1}$ -superficial element,
- (2) $x_i \in \mathcal{R}(\mathfrak{m})_1$ is $E_1(M_{i-1}) \oplus E_2(M_{i-1})$ -filter regular,

where $R_i = R/(x_1, \dots, x_i)$ and $M_i = M/(x_1, \dots, x_i)M$ for $i = 1, 2, \dots, d$. Set

$$J = (x_1, x_2, \dots, x_{d-1}), \quad S = R/J \quad \text{and} \quad N = M/JM.$$

Since $x_1, x_2, \dots, x_{d-1}$ is regular on R and M , ω and L are MCM , we get

$$\operatorname{Syz}_S^1(N) \cong \operatorname{Syz}_R^1(M)/J \operatorname{Syz}_R^1(M), \quad \omega_S = \omega_R/J\omega_R \quad \text{and} \quad \mu(M) = \mu(N).$$

By [7, 11(1)], we have

$$e_j(\omega_R) = e_j(\omega_S) \quad \text{and} \quad e_j(M) = e_j(N) \quad \text{for } j = 0, 1.$$

Since M and $\operatorname{Syz}_1^R(M)$ are MCM R -modules, by Corollary 3.4, we have

$$c_1(M) = c_1(N) \quad \text{and} \quad c_1(\operatorname{Syz}_R^1(M)) = c_1(\operatorname{Syz}_S^1(N)).$$

We now consider

$$\begin{aligned} c_1(N) + c_1(\operatorname{Syz}_S^1(N)) &= c_1(M) + c_1(\operatorname{Syz}_R^1(M)) \\ &= \mu(M) \cdot e_1(\omega_R), \quad \text{by our hypothesis} \\ &= \mu(M) \cdot e_1(\omega_S). \end{aligned}$$

Note that $\dim N = 1$. So, by induction hypothesis, N is free S -module. So $\operatorname{Ext}_S^1(N, k) = 0$. But $\operatorname{Ext}_R^{d+1}(M, k) \cong \operatorname{Ext}_S^1(N, k)$. This gives $\operatorname{Ext}_R^{d+1}(M, k) = 0$ and so $\beta_{d+1} = 0$. Therefore $\operatorname{projdim}_R(M) < \infty$. Since M is MCM , using Auslander-Buchsbaum's formula we get M is free. $\square$

5. THE CASE WHEN RING IS GORENSTEIN AND ITS ASSOCIATED GRADED RING IS COHEN-MACAULAY

In this section we assume that $(R, \mathfrak{m})$ is Gorenstein local ring of dimension d such that its associated graded ring $G_I(R)$ is CM . Let r be a reduction number of I . Then we prove

$$c_1^I(M) \geq r \cdot e_0^I(M) - \sum_{n=0}^{r-1} \sum_{j=0}^d \binom{d}{j} \ell \left(\text{Ext}_R^j(M, \frac{R}{I^{n+1-j}}) \right).$$

The above inequality motivates us to investigate the function

$$\Phi^I(M) = \sum_{j=0}^d \sum_{n=0}^{r-1} \binom{d}{j} \ell \left(\text{Ext}_R^j(M, \frac{R}{I^{n+1-j}}) \right).$$

This we do. We show that if $x \in I \setminus I^2$ is an R -superficial element then we have

$$\Phi^I(M) \geq \Phi^{I/(x)}(M/xM).$$

5.1. 0-Dimensional Case: We first deal with 0-dimensional Gorenstein ring S with unique maximal ideal $\mathfrak{n}$. Let J be a $\mathfrak{n}$ -primary ideal and N be a finitely generated S -module. Since S is 0-dimensional, there exists a positive integer $r \in \mathbb{N}$ such that $J^{r+1} = 0$ but $J^r \neq 0$. We have

$$\begin{aligned} \sum_{n=0}^{\infty} \ell \left(\text{Hom}_S(N, \frac{S}{J^{n+1}S}) \right) t^n &= \frac{f_N^J(t)}{1-t} \\ \sum_{n=0}^{r-1} \ell \left(\text{Hom}_S(N, \frac{S}{J^{n+1}S}) \right) t^n + \frac{\ell(\text{Hom}_S(N, S))}{1-t} \cdot t^r &= \frac{f_N^J(t)}{1-t} \end{aligned}$$

By Matlis duality, $\ell(\text{Hom}_S(N, S)) = \ell(N)$. So $c_0^J(N) = e_0^J(N)$. Thus

$$\begin{aligned} \sum_{n=0}^{r-1} \ell \left(\text{Hom}_S(N, \frac{S}{J^{n+1}S}) \right) t^n + \frac{e_0^J(N)}{1-t} \cdot t^r &= \frac{f_N^J(t)}{1-t} \\ \text{so, } f_N^J(t) &= (1-t) \cdot \sum_{n=0}^{r-1} \ell \left(\text{Hom}_S(N, \frac{S}{J^{n+1}S}) \right) t^n + e_0^J(N) \cdot t^r. \end{aligned}$$

Set $\alpha_n = \ell(\text{Hom}_S(N, S/J^{n+1}S))$. Therefore we get

$$(*) \quad f_N^J(t) = (1-t) \cdot \sum_{n=0}^{r-1} \alpha_n t^n + e_0^J(N) \cdot t^r.$$

Using (*) we compute $c_i^J(N)$ for $i = 0, 1$. Note that

$$\begin{aligned} c_0^J(N) &= f_N^J(1) = e_0^J(N) \\ \text{and } c_1^J(N) &= \left[\frac{d}{dt} f_N^J(t) \right]_{t=1} = r \cdot e_0^J(N) - \sum_{n=0}^{r-1} \alpha_n. \end{aligned}$$

5.2. General Case: Let $(R, \mathfrak{m})$ be Gorenstein local ring of dimension $d \geq 1$ with $\mathfrak{m}$ -primary ideal I such that the associated graded ring $G_I(R)$ is CM . We choose $\underline{x} = x_1, x_2, \dots, x_d \in I \setminus I^2$ be a sequence such that

- (1) x_i is I -superficial element with respect to R_{i-1} ,
- (2) $x_i \in \mathcal{R}(I)_1$ is $D_1(M_{i-1})$ -filter regular,

where $R_i = R/(x_1, \dots, x_i)$, $I_i = I/(x_1, \dots, x_i)$, $M_i = M/(x_1, \dots, x_i)M$ for $i = 1, 2, \dots, d$ and $D_1(M_i) = \bigoplus_{n \geq 0} \text{Ext}_{R_i}^1(M_i, R_i/I_i^n)$.

In the following proposition we set $S = R/(\underline{x})$, $\mathfrak{n} = \mathfrak{m}/(\underline{x})$, $J = I/(\underline{x})$ and $N = M/(\underline{x})M$.

Proposition 5.3. *(with hypotheses as in 5.2) We have*

$$c_1^I(M) \geq c_1^J(N).$$

Proof. By Corollary 3.4, we get

$$(i) \quad c_1^I(M) = c_1^{I_1}(M_1) = \dots = c_1^{I_{d-1}}(M_{d-1}).$$

So in view of (i), it is enough to show that

$$c_1^{I_{d-1}}(M_{d-1}) \geq c_1^J(N).$$

Thus we may assume that the dimension of ring is 1, i.e., $d = 1$. Since $G_I(R)$ is Cohen-Macaulay then R -superficial element $x \in I \setminus I^2$ implies that $x^* \in G_I(R)_1$ is $G_I(R)$ -regular. Set $S = R/(x)$, $N = M/xM$ and $J = I/(x)$. Thus we get a short exact sequence of the form

$$0 \rightarrow \frac{R}{I^n} \xrightarrow{\alpha_n^x} \frac{R}{I^{n+1}} \rightarrow \frac{S}{J^{n+1}} \rightarrow 0.$$

This gives the following long exact sequence

$$0 \rightarrow \text{Hom}_R(M, \frac{R}{I^n}) \rightarrow \text{Hom}_R(M, \frac{R}{I^{n+1}}) \rightarrow \text{Hom}_S(N, \frac{S}{J^{n+1}}) \rightarrow K_n \rightarrow 0,$$

where $K_n = \ker(\text{Ext}_R^1(M, R/I^n) \rightarrow \text{Ext}_R^1(M, R/I^{n+1}))$. By construction $K_n = 0$ for all $n \gg 0$. Therefore we have

$$\begin{aligned} \frac{f_R^M(t)}{(1-t)} + \sum_{n \geq 0} \ell(K_n)t^n &= \frac{f_S^N(t)}{(1-t)} \\ \Rightarrow f_S^N(t) &= f_R^M(t) + (1-t) \cdot \sum_{n \geq 0} \ell(K_n)t^n \end{aligned}$$

It follows that

$$\begin{aligned} c_1^J(N) &= c_1^I(M) - \sum_{n \geq 0} \ell(K_n) \\ \Rightarrow c_1^J(N) &\leq c_1^I(M). \end{aligned}$$

This completes the proof. $\square$

Before stating our main theorem we first prove the following lemma.

Lemma 5.4. *Let $x \in I \setminus I^2$ be an R -superficial element. Set $R' = R/(x)$, $I' = I/(x)$ and $M' = M/xM$ and . Then for $j \geq 0$ and $n \geq 1$ we have*

$$\begin{aligned} (i) \quad & \text{Ext}_{R'}^j(M', R'/I') = \text{Ext}_R^j(M, R/I) \\ (ii) \quad & \ell\left(\text{Ext}_{R'}^j(M', \frac{R'}{I'^{n+1}})\right) \leq \ell\left(\text{Ext}_R^j(M, \frac{R}{I^{n+1}})\right) + \ell\left(\text{Ext}_R^{j+1}(M, \frac{R}{I^n})\right). \end{aligned}$$

Proof. (i) The first assertion follows from [4, §18, Lemma 2].

(ii) Clearly $x \in I \setminus I^2$ is $G_I(R)$ -regular. Thus for each $n \geq 1$, we get an exact sequence of the form

$$0 \rightarrow \frac{R}{I^n} \xrightarrow{\alpha_n^x} \frac{R}{I^{n+1}} \rightarrow \frac{R'}{I'^{n+1}} \rightarrow 0.$$

Applying $\text{Hom}_R(M, -)$, we get following long exact sequence

$$\cdots \rightarrow \text{Ext}_R^j(M, \frac{R}{I^{n+1}}) \rightarrow \text{Ext}_R^j(M, \frac{R'}{I'^{n+1}}) \rightarrow \text{Ext}_R^{j+1}(M, \frac{R}{I^n}) \rightarrow \cdots$$

From [4, §18, Lemma 2] we get

$$\text{Ext}_R^j(M, \frac{R'}{I'^{n+1}}) = \text{Ext}_{R'}^j(M', \frac{R'}{I'^{n+1}})$$

The result follows. $\square$

We now make a convention that will be used throughout the section.

5.5. Convention: For ideal I , we set $I^j = R$ for $j \leq 0$.

A useful inequality is recorded in the following corollary.

Corollary 5.6. (with hypotheses as in 5.2) For all $n \geq 1$, we have

$$\ell\left(\text{Hom}_S(N, \frac{S}{J^n})\right) \leq \sum_{j=0}^d \binom{d}{j} \cdot \ell\left(\text{Ext}_R^j(M, \frac{R}{I^{n-j}})\right)$$

Proof. By repeated use of Lemma 5.4 we notice that

$$\begin{aligned} \ell\left(\text{Hom}_S(N, \frac{S}{J})\right) &= \ell\left(\text{Hom}_R(M, \frac{R}{I})\right) \\ \ell\left(\text{Hom}_S(N, \frac{S}{J^2})\right) &\leq \ell\left(\text{Hom}_R(M, \frac{R}{I^2})\right) + \binom{d}{1} \cdot \ell\left(\text{Ext}_R^1(M, \frac{R}{I})\right) \end{aligned}$$

Our convention gives that

$$\ell\left(\text{Hom}_S(N, \frac{S}{J^2})\right) \leq \sum_{j=0}^d \binom{d}{j} \cdot \ell\left(\text{Ext}_R^j(M, \frac{R}{I^{2-j}})\right).$$

Similarly one can check that for all $n \leq d$,

$$\ell\left(\text{Hom}_S(N, \frac{S}{J^n})\right) \leq \sum_{j=0}^d \binom{d}{j} \cdot \ell\left(\text{Ext}_R^j(M, \frac{R}{I^{n-j}})\right).$$

$\square$

A relation between the dual Hilbert coefficients $c_0^I(M)$ and $c_1^I(M)$ is established in the following theorem.

Theorem 5.7. (with hypotheses as in 5.2) Let r be the reduction number of I . Then

$$c_1^I(M) \geq r \cdot e_0^I(M) - \sum_{n=0}^{r-1} \sum_{j=0}^d \binom{d}{j} \ell\left(\text{Ext}_R^j(M, \frac{R}{I^{n+1-j}})\right)$$

Proof. By Proposition 5.3, we have $c_1^I(M) \geq c_1^J(N)$. Since $e_0^J(N) = e_0^I(M)$ then we get

$$c_1^I(M) \geq r \cdot e_0^I(M) - \sum_{n=0}^{r-1} \ell\left(\text{Hom}_S(N, \frac{S}{J^{n+1}})\right).$$

By Corollary 5.6, we have

$$\ell\left(\text{Hom}_S(N, \frac{S}{J^{n+1}})\right) \leq \sum_{j=0}^d \binom{d}{j} \cdot \ell\left(\text{Ext}_R^j(M, \frac{R}{I^{n+1-j}})\right)$$

Therefore

$$\sum_{n=0}^{r-1} \ell \left(\text{Hom}_S(N, \frac{S}{I^{n+1}}) \right) \leq \sum_{n=0}^{r-1} \sum_{j=0}^d \binom{d}{j} \ell \left(\text{Ext}_R^j(M, \frac{R}{I^{n+1-j}}) \right)$$

The result follows. $\square$

5.8. The previous result motivates us to study

$$\Phi^I(M) = \sum_{j=0}^d \sum_{n=0}^{r-1} \binom{d}{j} \ell \left(\text{Ext}_R^j(M, \frac{R}{I^{n+1-j}}) \right),$$

for a MCM R -module M of dimension d with respect to ideal I .

Remarks 5.9. We can simplify the expression for $\Phi^I(M)$.

(a) Notice that

$$\text{Ext}_R^j(M, \frac{R}{I^{n+1-j}}) = 0 \quad \text{for all } n < j.$$

Therefore we get

$$\Phi^I(M) = \sum_{j=0}^d \sum_{n=j}^{r-1} \binom{d}{j} \cdot \ell \left(\text{Ext}_R^j(M, \frac{R}{I^{n+1-j}}) \right)$$

(b) If $r - 1 \geq d$, then we have

$$\phi^I(M) = \sum_{j=0}^{r-1} \sum_{n=j}^{r-1} \binom{d}{j} \cdot \ell \left(\text{Ext}_R^j(M, \frac{R}{I^{n+1-j}}) \right)$$

Proposition 5.10. Let $x \in I \setminus I^2$ be an R -superficial element. Then we have

$$\Phi^I(M) \geq \Phi^{I/(x)}(M/xM)$$

Proof. For brevity set $R_1 = R/(x)$, $I_1 = I/(x)$ and $M_1 = M/xM$. Since $G_I(R)$ is Cohen-Macaulay and $x \in I \setminus I^2$ an R -superficial then we have an exact sequence of the form

$$0 \rightarrow \frac{R}{I^m} \xrightarrow{\alpha_m^x} \frac{R}{I^{m+1}} \rightarrow \frac{R_1}{I_1^{m+1}} \rightarrow 0.$$

This gives a long exact sequence of the form

$$(*) \quad \cdots \rightarrow \text{Ext}_R^j(M, \frac{R}{I^{m+1}}) \rightarrow \text{Ext}_{R_1}^j(M_1, \frac{R_1}{I_1^{m+1}}) \rightarrow \text{Ext}_R^{j+1}(M, \frac{R}{I^m}) \rightarrow \cdots$$

By definition of Φ , as $\dim M_1 = d - 1$, we have

$$\Phi^{I_1}(M_1) = \sum_{j=0}^{d-1} \sum_{n=0}^{r-1} \binom{d-1}{j} \ell \left(\text{Ext}_{R_1}^j(M_1, \frac{R_1}{I_1^{n+1-j}}) \right)$$

From (*), it follows that

$$\begin{aligned}
\Phi^{I_1}(M_1) &\leq \sum_{j=0}^{d-1} \sum_{n=0}^{r-1} \binom{d-1}{j} \ell \left(\text{Ext}_R^j(M, \frac{R}{I^{n+1-j}}) \right) + \\
&\quad \sum_{j=0}^{d-1} \sum_{n=0}^{r-1} \binom{d-1}{j} \ell \left(\text{Ext}_R^{j+1}(M, \frac{R}{I^{n-j}}) \right) \\
&= \sum_{j=0}^{d-1} \sum_{n=0}^{r-1} \binom{d-1}{j} \ell \left(\text{Ext}_R^j(M, \frac{R}{I^{n+1-j}}) \right) + \\
&\quad \sum_{j=1}^d \sum_{n=0}^{r-1} \binom{d-1}{j-1} \ell \left(\text{Ext}_R^j(M, \frac{R}{I^{n+1-j}}) \right).
\end{aligned}$$

Notice that

$$\binom{d-1}{j} = 0 \quad \text{for } j = d \text{ and } \binom{d-1}{j-1} = 0 \quad \text{for } j = 0.$$

Therefore using a well-known Binomial identity we get

$$\Phi^{I_1}(M_1) \leq \sum_{j=0}^d \sum_{n=0}^{r-1} \binom{d}{j} \ell \left(\text{Ext}_R^j(M, \frac{R}{I^{n+1-j}}) \right).$$

Hence it follows that $\Phi^{I_1}(M_1) \leq \Phi^I(M)$. $\square$

The following is an immediate consequence.

Corollary 5.11. *Let the situation be as in 5.2. Then we have*

$$\Phi(M) \geq \Phi(M_1) \geq \Phi(M_2) \geq \cdots \geq \Phi(M_d) = \Phi(N).$$

Remark 5.12. Let S be 0-dimensional Gorenstein local ring, and assume that $J^{r+1} = 0$ but $J^r \neq 0$. Then

$$\Phi^J(N) = \sum_{n=0}^{r-1} \ell \left(\text{Hom}_S(N, \frac{S}{J^{n+1}}) \right)$$

Example 5.13. Let $R = \mathbf{Q}[x, y]/(f)$ be a hypersurface ring of dimension $d = 1$, where $f = x^2 + xy + y^2$ a homogeneous polynomial of degree 2. Let $\mathfrak{m} = \langle x, y \rangle$ R be a maximal ideal of R . Consider an R -module

$$M = \left\langle \begin{pmatrix} x \\ -y \end{pmatrix}, \begin{pmatrix} x+y \\ x \end{pmatrix} \right\rangle \subseteq R^2.$$

Note that M is an *MCM* R -module. Clearly reduction number r of $\mathfrak{m}$ is 1. Therefore we get

$$\begin{aligned}
\Phi^{\mathfrak{m}}(M) &= \sum_{j=0}^d \sum_{n=0}^{r-1} \binom{d}{j} \ell \left(\text{Ext}_R^j(M, \frac{R}{\mathfrak{m}^{n+1-j}}) \right) \\
&= \ell \left(\text{Hom}_R(M, \frac{R}{\mathfrak{m}}) \right) \\
&= \mu(M) = 2.
\end{aligned}$$

$$\Rightarrow r \cdot e_0(M) - \Phi^{\mathfrak{m}}(M) = 0.$$

Therefore we get $c_1^{\mathfrak{m}}(M) \geq 0$.

We now give an example where $c_1^I(M)$ could be negative.

Example 5.14. Let $(R, \mathfrak{m})$ be zero dimensional Gorenstein local ring with reduction number of $\mathfrak{m} = 2$, i.e., $\mathfrak{m}^2 \neq 0$ but $\mathfrak{m}^3 = 0$. Then we get

$$(i) \quad \mathfrak{m}^2 = \text{Soc}(R) = \text{Hom}_R(k, R) \simeq k.$$

We now take $M = k$. Consider the exact sequence

$$0 \longrightarrow \mathfrak{m}^2 \longrightarrow R \longrightarrow \frac{R}{\mathfrak{m}^2} \longrightarrow 0.$$

This induces the following isomorphism

$$(ii) \quad \text{Hom}_R(k, \frac{R}{\mathfrak{m}^2}) \simeq \text{Ext}_R^1(k, k).$$

Let $\mu(\mathfrak{m}) = h$, the minimal number of generators of $\mathfrak{m}$. Then using (i) and (ii), we get

$$\begin{aligned} \sum_{n \geq 0} \ell \left(\text{Hom}_R(k, \frac{R}{\mathfrak{m}^{n+1}}) \right) t^n &= \frac{f_M^{\mathfrak{m}}(t)}{1-t} \\ \Rightarrow 1 + ht + \frac{t^2}{1-t} &= \frac{f_M^{\mathfrak{m}}(t)}{1-t}. \\ \text{So, } f_M^{\mathfrak{m}}(t) &= t^2 + (1-t)(1+ht). \end{aligned}$$

Therefore it follows that $c_1^{\mathfrak{m}}(M) = 1 - h$. So $c_1^{\mathfrak{m}}(M) < 0$ when $h \geq 2$.

A specific example of the above kind is the following:

Example 5.15. Let e be any positive integer greater than 3. Consider

$$R = \frac{k[[t^e, t^{e+1}, \dots, t^{2e-2}]]}{(t^e)}$$

It follows from [8, 3.2] that R is 0-dimensional Gorenstein local ring with maximal ideal

$$\mathfrak{m} = \frac{(t^e, t^{e+1}, \dots, t^{2e-2})}{(t^e)}.$$

Clearly $h = \mu(\mathfrak{m}) = e - 2 \geq 2$.

6. EXAMPLE: DUAL HILBERT-SAMUEL FUNCTION OF AN ULRICH MODULE

In this section we assume that R and its associated graded ring $G_{\mathfrak{m}}(R)$ are Gorenstein. Let $L = \text{Syz}_R^1(M)$, denote the first syzygy module of M . Set $S = R/(x)$, $N = M/xM$ and $\mathfrak{n} = \mathfrak{m}/xR$. The goal of this section is to compute the Dual Hilbert-Samuel function of an Ulrich module. Recall that a MCM module M is said to be Ulrich if $e_0^{\mathfrak{m}}(M) = \mu(M)$.

The following theorem gives a sufficient condition which ensures that the dual Hilbert-Samuel function of M behaves "perfectly" with respect to a superficial element.

Theorem 6.1. *Assume that $x \in \mathfrak{m} \setminus \mathfrak{m}^2$ is such that x^* is $G_{\mathfrak{m}}(R) \oplus G_{\mathfrak{m}}(L)$ -regular. Then*

$$\varepsilon_M^0(\mathfrak{m}, n) - \varepsilon_M^0(\mathfrak{m}, n-1) = \varepsilon_N^0(\mathfrak{n}, n) \quad \text{for } n \geq 0.$$

Proof. Consider the following exact sequence

$$0 \rightarrow \mathfrak{m}^n \rightarrow R \rightarrow \frac{R}{\mathfrak{m}^n} \rightarrow 0.$$

This induces that

$$(i) \quad D_{\mathfrak{m}}^1(M) \simeq \bigoplus_{n \geq 0} \text{Ext}_R^2(M, \mathfrak{m}^n) \simeq \text{Ext}_R^2(M, \mathcal{R}(\mathfrak{m})).$$

Also the exact sequence

$$0 \rightarrow L \rightarrow R^{\mu(M)} \rightarrow M \rightarrow 0$$

yields the following isomorphism

$$(ii) \quad \text{Ext}_R^1(L, \mathcal{R}(\mathfrak{m})) \simeq \text{Ext}_R^2(M, \mathcal{R}(\mathfrak{m})).$$

From (i) and (ii), it follows that

$$(iii) \quad D_{\mathfrak{m}}^1(M) \simeq \text{Ext}_R^1(L, \mathcal{R}(\mathfrak{m})).$$

Since $G_{\mathfrak{m}}(R)$ is Gorenstein and $G_{\mathfrak{m}}(L)$ and the homogeneous element $x^* \in \mathcal{R}(\mathfrak{m})_1$ is $G_{\mathfrak{m}}(R) \oplus G_{\mathfrak{m}}(L)$ -regular then by [6, 11.5], the sequence

$$0 \rightarrow \text{Hom}_R(L, \mathfrak{m}^n) \rightarrow \text{Hom}_R(L, \mathfrak{m}^{n+1}) \rightarrow \text{Hom}_R(L, \mathfrak{n}^{n+1}) \rightarrow 0$$

is exact for all $n \geq 0$. Therefore it follows that the maps

$$\text{Ext}_R^1(L, \mathfrak{m}^n) \rightarrow \text{Ext}_R^1(L, \mathfrak{m}^{n+1}) \quad \text{are injective for all } n \geq 0$$

Using (iii) the maps

$$\text{Ext}_R^1(M, \frac{R}{\mathfrak{m}^n}) \xrightarrow{xt} \text{Ext}_R^1(M, \frac{R}{\mathfrak{m}^{n+1}})$$

are injective for all $n \geq 1$. Therefore the short exact sequence

$$0 \rightarrow \frac{R}{\mathfrak{m}^n} \xrightarrow{\alpha_n^x} \frac{R}{\mathfrak{m}^{n+1}} \rightarrow \frac{S}{\mathfrak{n}^{n+1}} \rightarrow 0.$$

induces a short exact of the form

$$0 \rightarrow \text{Hom}_R(M, \frac{R}{\mathfrak{m}^n}) \xrightarrow{x} \text{Hom}_R(M, \frac{R}{\mathfrak{m}^{n+1}}) \rightarrow \text{Hom}_R(N, \frac{S}{\mathfrak{n}^{n+1}}) \rightarrow 0.$$

This proves our assertion. $\square$

6.2. The case when M is Ulrich module: Let M be Ulrich R -module. Then there exists $J = (x_1, x_2, \dots, x_d)$, a minimal reduction of $\mathfrak{m}$ such that $JM = \mathfrak{m}M$. Therefore we get

$$G_{\mathfrak{m}}(M) \simeq \frac{M}{\mathfrak{m}M}[\bar{x}_1, \bar{x}_2, \dots, \bar{x}_d].$$

Let $\mu(M)$ denote minimal number of generators of M . Then $M/JM \simeq k^{\mu(M)}$.

6.3. Assume $(S, \mathfrak{n}, k)$ is a 0-dimensional Gorenstein local ring with $G_n(S)$ Gorenstein. So there exists a positive integer r such that $\mathfrak{n}^r \neq 0$ but $\mathfrak{n}^{r+1} = 0$. From the exact sequence

$$0 \rightarrow \mathfrak{n}^i \rightarrow S \rightarrow \frac{S}{\mathfrak{n}^i} \rightarrow 0$$

one can verify that

$$\text{Hom}_S(k, \frac{S}{\mathfrak{n}^i}) \simeq \begin{cases} \text{Ext}_S^1(k, \mathfrak{n}^i) & \text{for } i = 1, 2, \dots, r \\ k & \text{for } i > r. \end{cases}$$

Also notice that $\text{Ext}_S^1(k, \mathfrak{n}^i) = k^{\mu_1(\mathfrak{n}^i)}$, where $\mu_1(\mathfrak{n}^i)$ is the first *Bass number* of $\mathfrak{n}^i$. Now we can compute the dual Hilbert-Samuel series of k with respect to $\mathfrak{n}$.

$$\begin{aligned} D(k, t) &= \mu_1(\mathfrak{n}) + \mu_1(\mathfrak{n}^2)t + \dots + \mu_1(\mathfrak{n}^r)t^{r-1} + \frac{t^{r+1}}{(1-t)} \\ &= \frac{(1-t) \cdot [\sum_{i=0}^{r-1} \mu_1(\mathfrak{n}^{i+1}) \cdot t^i] + t^{r+1}}{(1-t)} \end{aligned}$$

Proposition 6.4. *Let M be an Ulrich R -module. Then we have*

$$D(M, t) = \frac{\mu(M) \cdot [(1-t) \cdot [\sum_{i=0}^{r-1} \mu_1(\mathfrak{n}^{i+1}) \cdot t^i] + t^{r+1}]}{(1-t)^{d+1}}.$$

Proof. Since M is Ulrich R -module, there exists $J = (x_1, x_2, \dots, x_d)$, a minimal reduction of $\mathfrak{m}$ such that $JM = \mathfrak{m}M$. Set $\bar{M} = M/JM$. In [5] we prove that $\text{Gm}(\text{Syz}_1^A(M))$ is Cohen-Macaulay. In view of Theorem 6.1, it follows that

$$\begin{aligned} D(M, t) &= \frac{D(\bar{M}, t)}{(1-t)^d} = \frac{D(k^{\mu(M)}, t)}{(1-t)^d} \\ &= \mu(M) \cdot \frac{D(k, t)}{(1-t)^d} \\ &= \frac{\mu(M) \cdot [(1-t) \cdot [\sum_{i=0}^{r-1} \mu_1(\mathfrak{n}^{i+1}) \cdot t^i] + t^{r+1}]}{(1-t)^{d+1}} \end{aligned}$$

The last equality follows from 6.3. □

REFERENCES

- [1] M. F. Atiyah and I. G. Macdonald, *Introduction to commutative algebra*, Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1969. MR MR0242802
- [2] M.P. Brodmann and R.Y. Sharp, *Local cohomology : an algebraic introduction with geometric applications*, Cambridge Studies in Advanced Mathematics, vol. 60, Cambridge University Press, Cambridge, 1998.
- [3] W. Bruns and J. Herzog, *Cohen-Macaulay rings*, Cambridge Studies in Advanced Mathematics, vol. 39, Cambridge University Press, Cambridge, 1993. MR MR1251956
- [4] H. Matsumura, *Commutative ring theory*, Cambridge Studies in Advanced Mathematics, vol. 8, Cambridge University Press, Cambridge, 1986, Translated from the Japanese by M. Reid. MR MR879273
- [5] T. J. Puthenpurakal, *Dual filtrations and applications*, work in progress.
- [6] ———, *Gorenstein approximation, dual filtrations and applications*, Preprint. arXiv:0807.0471.
- [7] ———, *Hilbert-coefficients of a Cohen-Macaulay module*, J. Algebra **264** (2003), no. 1, 82–97. MR MR1980687
- [8] J. D. Sally, *Tangent cones at Gorenstein singularities*, Compositio Math. **40** (1980), no. 2, 167–175. MR MR563540
- [9] E. Theodorescu, *Derived functors and Hilbert polynomials*, Math. Proc. Cambridge Philos. Soc. **132** (2002), no. 1, 75–88. MR MR1866325

DEPARTMENT OF MATHEMATICS, IIT BOMBAY, POWAI, MUMBAI 400 076, INDIA
E-mail address: tputhen@math.iitb.ac.in

DEPARTMENT OF MATHEMATICS, IIT BOMBAY, POWAI, MUMBAI 400 076, INDIA